

# **Bulletin**

of the

# **Chicago Academy of Sciences**

**The Influence of the Balcones Escarpment on  
the Distribution of Amphibians  
and Reptiles in Texas**

**HOBART M. SMITH AND HELMUT K. BUECHNER**



**Chicago**  
*Published by the Academy*  
**1947**

*The Bulletin of the Chicago Academy of Sciences* was initiated in 1883 and volumes 1 to 4 were published prior to June, 1913. During the following twenty-year period it was not issued. Volumes 1, 2, and 4 contain technical or semi-technical papers on various subjects in the natural sciences. Volume 3 contains museum reports, descriptions of museum exhibits, and announcements.

Publication of the *Bulletin* was resumed in 1934 with volume 5 in the present format. It is now regarded as an outlet for short to moderate-sized original papers on natural history, in its broad sense, by members of the museum staff, members of the Academy, and for papers by other authors which are based in considerable part upon the collections of the Academy. It is edited by the Director of the Museum with the assistance of a committee from the Board of Scientific Governors. The separate numbers are issued at irregular intervals and distributed to libraries and scientific organizations, and to specialists with whom the Academy maintains exchanges. A reserve is set aside for future need as exchanges and the remainder of the edition offered for sale at a nominal price. When a sufficient number of pages have been printed to form a volume of convenient size, a title page, table of contents, and index are supplied to libraries and institutions which receive the entire series.

Howard K. Gloyd, Director of the Museum

*Committee on Publications:*

Alfred Emerson, Professor of Zoology, University of Chicago.  
C. L. Turner, Professor of Zoology, Northwestern University.  
Hanford Tiffany, Professor of Botany, Northwestern University.

# Bulletin of the Chicago Academy of Sciences

---

## The Influence of the Balcones Escarpment on the Distribution of Amphibians and Reptiles in Texas

HOBART M. SMITH AND HELMUT K. BUECHNER\*

It is a common observation that the geographic ranges of many plants and animals are limited by the Balcones Escarpment in Texas, but little critical study has been made of the role of the escarpment as a natural boundary. The area offers much material for studies in speciation. The escarpment is in fact the eastern edge of the Edwards Plateau, an elevated area of features markedly different from those of adjacent territories. The plateau is xeric in nature, having less rainfall, scrubbiier and more scant vegetation, a porous soil, little standing water, and an evaporation rate exceeding the precipitation rate. Surrounding lowlands are of mesic character, with more rainfall, lush vegetation, less porous soil, much standing water, and a rainfall rate exceeding the evaporation rate.

Various degrees and sorts of distributional limitation are exhibited by the numerous species occurring there, depending upon interaction of the wide variety of ecological conditions on the one hand, with on the other hand the physiological tolerances, life histories, habits, evolutionary history, etc., of the individual species. As the conditions are limited in occurrence, so are the animals limited; and as there are innumerable combinations of potentially limiting factors, so are there innumerable distributional patterns among the species concerned. The purpose of the present paper is to analyze the extent of limitation exercised by the factors correlated with the escarpment upon the geographical distribution of reptiles and amphibians in Texas.

Most of the material for the discussion presented here was plotted in map form for all the amphibians and reptiles reported from Texas. These maps are based solely upon published reports, materials in the Texas Cooperative Wildlife Collection at the Agricultural and Mechanical College of Texas, and specimens in the private collection of Bryce C. Brown.




Figure 1. The Edwards Plateau of Texas, indicated by stippled area. The Balcones Escarpment forms its eastern and southern boundaries.

#### FEATURES OF THE ESCARPMENT AND ADJACENT AREAS

*Location.* The Balcones Escarpment forms an arc through central Texas, starting from Waco, extending southward through Austin to San Antonio, and then swinging west to Del Rio on the Rio Grande (see Fig. 1). It is not a single long vertical cliff, but rather a series of breaks in the earth's crust, forming cliffs, hills and plateaus up to a height of about 1000 feet. The highest parts are in the vicinity of Austin, San Antonio, and Uvalde; northward from Waco and westward from Uvalde the escarpment gradually decreases in height until it is recognizable only through the presence of small fault lines.

*Climate.* In passing through a distance of 10 to 20 miles across and at right angles to the escarpment in the vicinity of Austin, San Antonio, or Uvalde, one encounters conspicuous changes in the climate, vegetation, and animal life. In many places the change in vegetation is as sharply defined at the fault lines as a fence row between areas under different types of land use.

The Edwards Plateau, which lies northwest of the escarpment, has a much drier climate than the area to the east of the plateau. The average rainfall is about 25 to 30 inches on the plateau, about 10 to 15 inches less than the area east of the escarpment. The plateau temperatures are a few degrees cooler than the lower elevations. During every month of the year the evaporation on the plateau exceeds the amount of precipitation. This accounts for the dryness of the climate. Very little of the water which falls is available for plant growth, and permanent bodies of open water are seldom found. The fault area of the escarpment, however, is characterized by many clear, spring-fed streams. There are also several large caves with permanent water.

*Vegetation.* The xeric nature of the biota west of the escarpment is well expressed in the plateau vegetation (Buechner, 1944). An extensive carpet of oak and cedar covers most of the area, with eastern mesophytes (cypress, elm, hickory, ash, birch) finding a suitable habitat along the main streams. The oak-cedar woodland is more closely related to the vegetation of the lower slopes of the Rocky Mountains than to the blackland prairie and post oak-hickory forest to the east or the mesquite chaparral to the south (Tharp, 1939). There is a conspicuous difference not only in species of plants, but also in the growth form of the vegetation. As a result of the dry climate, the tree species are stunted, forming a midget woodland, which becomes a low mat at higher elevations on hill tops as well as farther west on the plateau.

*Animal Life.* The influence of the escarpment on animal life is as marked as the correlation between climate, soils, and vegetation. This is to be expected, since animal life, vegetation, and the physical environment all interact to produce the large entities known as ecosystems. Specific comparisons of the mammalian fauna and avifauna on either side of the escarpment have never been made. Nevertheless, in comparing the birds of Kerr County on the Edwards Plateau with those of Brazos County about 150 miles to the east (Buechner, 1946), it becomes obvious that the Balcones Escarpment has a definite limiting influence on birds. The birds of Kerr County have greater affinities to the more western bird life than to that of the east. The breeding birds there consist of more than twice as many western and southwestern forms as eastern. About 56 per cent of all the breeding birds in Kerr County find their range limits in the county, which is not far from the escarpment.

This figure compares fairly well with the 77 per cent of amphibians and reptiles that find their range limits at the escarpment.

A further study of other animals and of plants involved at the escarpment may lead to a definite percentage criterion useful in designating biotic areas. Indications from the present study are that the amount of permissible overlap at a natural boundary may be much higher than normally expected, even when a large number of plant and animal species are brought into consideration.

#### INFLUENCE OF THE ESCARPMENT ON AMPHIBIANS AND REPTILES

Since there is considerable variance in the effect of the escarpment on the various groups of amphibians and reptiles, our analysis is made by natural groupings. The species and subspecies were segregated, on the basis of their distribution, into several categories. We tabulated them, for each of the five groups (to wit, salamanders, anurans, lizards, snakes, turtles), as follows: (1) limited (that is to say, kinds which occur at the escarpment but do not cross it extensively in any direction) classified in turn as eastern, western, northern, southern or endemic (autochthonous species, defined as species originating in a certain area but now occurring to a limited extent in adjacent areas, were counted as endemic); (2) overlapping (kinds which occur extensively on both sides of the escarpment); or (3) peripheral (kinds which do not occur at the escarpment). The latter we again grouped as eastern, western, southern or northern.

We found good evidence of the existence in the state of 38 kinds of anurans, 18 salamanders, 44 lizards, 93 snakes, and 23 turtles, making a total of 216. Fifty-six per cent (122) of the total Texan herpetofauna (with a range of from 50 to 67 per cent among the 5 groups) reach the escarpment. Of these, 77 per cent (94) are limited there, and 11 per cent (total range 1 to 42 per cent) are endemic.

Not all of the range limits considered here are sharply marked off. Among those designated as limited by the escarpment, there may be a slight overstepping, particularly at the Del Rio and Waco ends where the escarpment levels out. In a number of cases the eastern portion of the escarpment forms a sharply defined boundary of the range of the species or subspecies while the southern face is insignificant and the range extends southward over it or beyond the entire southern portion of the Rio Grande. In all such cases it is obvious that low humidity, which occurs in an area similarly outlined, is more significant than any other factor. Species thus partially limited (i.e., by only the eastern face of the plateau) we have nevertheless tabulated as "limited."

It is obvious that no two independent analyses by different investigators are likely to arrive at exactly the same conclusion, since estimates of poorly known ranges must at the present time be highly arbitrary. Likewise arbitrary is the extent of overlapping of the escarpment accepted as commensurate with the definition of "limited" forms. These and other variables make the present estimates subject to later revision.

*Anurans.* There are 38 forms of the order Salientia whose reported occurrence in Texas is unquestioned. Of these, 50 per cent reach the escarpment. About half of the remaining 50 per cent are of eastern distribution, and most of these belong to the genera *Hyla* and *Rana*, whose requirements for a humid climate are apparently so restricted that they are unable to range as far west as the escarpment.

Of the frogs and toads that do reach the Balcones region, 74 per cent find their range limits there. Most of these are eastern frogs whose requirements for an aquatic habitat cannot be met extensively enough beyond the escarpment. Western forms which find their eastern limit there are toads and spadefoots adapted to a xeric type of habitat. Two frogs, *Eleutherodactylus latrans* and *Syrhophus marnockii*, are adapted to the limestone ledges, cliffs, and caves that front the Edwards Plateau, and are endemic to this area. The species may be tabulated as follows.

#### LIMITED BY ESCARPMENT

##### Western

*Bufo cognatus*  
*Bufo debilis*  
*Bufo punctatus*  
*Scaphiopus couchii*  
*Scaphiopus hammondi*

##### Eastern

*Bufo valliceps valliceps*  
*Hyla cinerea cinerea*  
*Hyla versicolor chrysoscelis*  
*Pseudacris clarkii*  
*Pseudacris streckeri*  
*Rana catesbeiana*  
*Scaphiopus hurteri*

##### Endemic

*Eleutherodactylus latrans*  
*Syrhophus marnockii*

#### OCCURRING ON BOTH SIDES

*Acris crepitans*  
*Bufo compactilis*  
*Bufo woodhousii woodhousii* (mostly west)  
*Microhyla olivacea*  
*Rana pipiens*

# NOT REACHING ESCARPMENT

## Western

*Bufo insidiosus*  
*Hyla arenicolor*  
*Scaphiopus bombifrons*  
*Syrrophus gaigeae*

## Southern

*Bufo horribilis*  
*Hyla baudinii baudinii*  
*Hypopachus cuneus cuneus*  
*Leptodactylus labialis*  
*Syrrophus campi*

## Eastern

*Bufo woodhousii fowleri*  
*Hyla crucifer crucifer*  
*Hyla femoralis*  
*Hyla squirella*  
*Hyla versicolor versicolor*  
*Pseudacris nigrita triseriata*  
*Microhyla carolinensis*  
*Rana areolata areolata*  
*Rana clamitans*  
*Rana palustris*

*Salamanders.* Of the 18 kinds of salamanders whose reported occurrence has been verified, 67 per cent occur at the escarpment, and all of these are at their range limits. The five kinds endemic (or autochthonous) to the escarpment are correlated in distribution with the caves and cave waters, artesian wells, and spring-fed streams of the area. All but one of the non-endemic salamanders are eastern or southern forms. Since this group of amphibians is more restricted to an aquatic environment than is any other group under consideration, it is the one affected most by the change towards drier conditions at the Balcones region. The species may be tabulated as follows.

# LIMITED BY ESCARPMENT

## Western

*Ambystoma tigrinum mavortium*

## Southern

*Diemictylus meridionalis*

## Endemic

*Eurycea latitans*  
*Eurycea nana*  
*Eurycea neotenes*  
*Plethodon glutinosus albagula*  
*Typhlomolge rathbuni*

## Eastern

*Ambystoma texanum*  
*Ambystoma tigrinum tigrinum*  
*Manculus quadridigitatus*  
*Plethodon glutinosus glutinosus*  
*Siren intermedia nettingi*

# NOT REACHING ESCARPMENT

## Eastern

*Ambystoma maculatum*  
*Ambystoma opacum*  
*Ambystoma talpoideum*  
*Amphiuma means tridactylum*  
*Desmognathus fuscus brimleyorum*  
*Diemictylus viridescens louisianensis*



*Lizards.* Forty-four lizards are recorded in Texas, and 52 per cent of them reach the escarpment. Of these, all but two (95 per cent) are limited at the fault area. In contrast to amphibians, which require water for at least a stage of their life history, the reptiles are much more tolerant of the dry conditions on the Edwards Plateau: there are more western species limited by the escarpment than there are of eastern and southern species. The reverse is true of amphibians. The species of lizards are as follows.

#### LIMITED BY ESCARPMENT

##### Western

*Cnemidophorus gularis gularis*  
*Coleonyx brevis*  
*Crotaphytus collaris collaris*  
*Eumeces brevilineatus*  
*Eumeces obsoletus*  
*Gerrhonotus liocephalus infernalis*  
*Holbrookia texana*  
*Sceloporus poinsettii*  
*Sceloporus undulatus consobrinus*  
*Urosaurus ornatus ornatus*

##### Southern

*Eumeces tetragrammus*  
*Sceloporus variabilis marmoratus*

##### Eastern

*Anolis carolinensis*  
*Cnemidophorus sexlineatus*  
*Eumeces fasciatus*  
*Eumeces laticeps*  
*Holbrookia propinqua*  
*Leiopisma laterale*  
*Ophisaurus ventralis*  
*Sceloporus undulatus hyacinthinus*

##### Endemic

*Holbrookia maculata lacerata*

#### OCCURRING ON BOTH SIDES

*Phrynosoma cornutum*  
*Sceloporus olivaceus*

#### NOT REACHING ESCARPMENT

##### Western

*Cnemidophorus grahamii*  
*Cnemidophorus gularis octolineatus*  
*Cnemidophorus perplexus*  
*Cnemidophorus tessellatus tessellatus*  
*Crotaphytus collaris baileyi*  
*Phrynosoma douglassii hemandesi*  
*Phrynosoma modestum*  
*Sceloporus magister magister*  
*Sceloporus merriami merriami*

*Eumeces gagei*  
*Eumeces multivirgatus*  
*Eumeces taylori*  
*Gambelia wislizenii wislizenii*  
*Holbrookia maculata approximans*  
*Sceloporus merriami annulatus*  
*Urosaurus ornatus schmidtii*  
*Uta stansburiana stejnegeri*

##### Eastern

*Eumeces anthracinus*

##### Southern

*Crotaphytus reticulatus*  
*Sceloporus grammicus disparilis*  
*Sceloporus cyanogenys*

**Snakes.** Of the 93 species or subspecies of snakes for which we find authentic records from Texas, 53 range up to or across the escarpment; 57 per cent of the ophidian fauna is thus involved in the escarpment area. Of those involved, 70 per cent are limited by the escarpment. Most of the forms not reaching the cliffs are western species. Only ten are eastern species, including three forms of *Natrix* and one of *Farancia*. The species of snakes which are limited are divided almost equally among mesic eastern forms and xeric western or southern forms, indicating a wide range in physiological adaptivity—wider by all odds than in any other reptilian or amphibian order or suborder. The species may be tabulated as follows.

#### LIMITED BY ESCARPMENT

##### Western

*Agkistrodon mokeson laticinctus*  
*A. rizona elegans elegans*  
*Crotalus molossus molossus*  
*Crotalus*  
*Diadophis punctatus docilis*  
*Gyalopion canum*  
*Hypsiglena ochrorhyncha texana*  
*Lampropeltis getulus splendida*  
*Lampropeltis triangulum gentilis*  
*Leptotyphlops dulcis*  
*Masticophis taeniatus omatus*  
*Rhinocheilus lecontei tessellatus*  
*Sonora episcopa*  
*Thamnophis eques cyrtopsis*

##### Eastern

*Agkistrodon mokeson mokeson*  
*Agkistrodon piscivorus leucostomus*  
*Crotalus horridus atricaudatus*  
*Diadophis punctatus stictogenys*  
*Haldea striatula*  
*Haldea valeriae elegans*  
*Heterodon contortrix contortrix*  
*Lampropeltis getulus holbrooki*  
*Lampropeltis triangulum amaura*  
*Micrurus fulvius tenere*  
*Natrix grahamii*  
*Natrix sipedon confiuens*  
*Sistrurus miliarius streckeri*  
*Troidoclonion lineatum*  
*Tantilla gracilis*  
*Storeria occipitomaculata occipitomaculata*  
*Thamnophis sirtalis sirtalis*

##### Southern

*Coluber constrictor stejnegerianus*  
*Masticophis taeniatus schotti*  
*Sonora taylori*  
*Tantilla nigriceps fumiceps*

##### Endemic

*Natrix harteri*  
*Thamnophis rufipunctatus (in this area)*

#### OCCURRING ON BOTH SIDES

*Coluber constrictor flaviventris*  
*Crotalus atrox*  
*Elaphe laeta laeta*  
*Elaphe obsoleta confinis*  
*Heterodon nasicus nasicus*  
*Lampropeltis calligaster calligaster*  
*Masticophis flagellum testaceus*  
*Natrix erythrogaster transversa*

*Natrix rhombifera rhombifera*  
*Opheodrys aestivus*  
*Pituophis catenifer sayi*  
*Salvadora lineata*  
*Sistrurus catenatus tergeminus*  
*Storeria dekayi texana*  
*Thamnophis marciana*  
*Thamnophis sauritus proximus*

# NOT REACHING ESCARPMENT

## Western

*Agkistrodon mokeson pictigaster*  
*Crotalus lepidus lepidus*  
*Crotalus lepidus klauberi*  
*Crotalus scutulatus scutulatus*  
*Diadophis regalis regalis*  
*Elaphe bairdi*  
*Elaphe subocularis*  
*Hypsiglena ochrorhyncha ochrorhyncha*  
*Lampropeltis alterna*  
*Leptotyphlops humilis segregus*  
*Leptotyphlops myopica dissecta*  
*Masticophis taeniatus taeniatus*  
*Salvadora grahamiae*  
*Salvadora hexalepis deserticola*  
*Sonora semiannulata blanchardi*  
*Tantilla atriceps*  
*Tantilla nigriceps nigriceps*  
*Thamnophis macrostemma megalops*  
*Thamnophis sirtalis parietalis*  
*Trimorphodon wilkinsonii*

## Northern

*Diadophis punctatus amnyi*  
*Opheodrys vernalis blanchardi*

## Eastern

*Agkistrodon mokeson austrinus*  
*Carphophis amoenus vermis*  
*Coluber constrictor constrictor*  
*Crotalus horridus horridus*  
*Farancia abacura reinwardtii*  
*Masticophis flagellum flagellum*  
*Natrix cyclopion cyclopion*  
*Natrix erythrogaster erythrogaster*  
*Natrix sipedon clarkii*  
*Rhadinaea flavilata*

## Southern

*Coniophanes imperialis imperialis*  
*Dryomarchon corais erebennus*  
*Dryomobius margaritiferus margaritiferus*  
*Ficimia streckeri*  
*Heterodon nasicus kennerlyi*  
*Lampropeltis triangulum annulata*  
*Leptodeira annulata septentrionalis*  
*Masticophis taeniatus ruthveni*

**Turtles.** The turtles are considerably less tolerant of environmental conditions than other reptiles. There are 23 species of freshwater turtles authentically reported in Texas, 15 of which reach the faulted area. Ten of the latter, or 67 per cent, are restricted. All but one of these are eastern or southern species. Perhaps one reason for the relatively high percentage of range limitations among the turtles is that proper egg-laying conditions are not found on the Edwards Plateau. The soil is so thin and rocky on the plateau that only a small percentage of the area provides soil of sufficient depth and proper texture for burying eggs. The species may be listed as follows.

# LIMITED BY ESCARPMENT

## Eastern

*Amyda mutica*  
*Chelydra serpentina serpentina*  
*Graptemys pseudographica kohnii*  
*Kinostemon subrubrum hippocrepis*  
*Macrochelys temminckii*  
*Sternotherus carinatus*  
*Sternotherus odoratus*  
*Terrapene carolina triunguis*

## Southern

*Gopherus berlandieri*

## Endemic

*Graptemys pseudogeographica versa*

#### OCCURRING ON BOTH SIDES

*Amyda emoryi*  
*Kinosternon flavescens*  
*Pseudemys floridana texana*  
*Pseudemys scripta elegans*  
*Terrapene ornata*

#### NOT REACHING ESCARPMENT

##### Western

*Chrysemys pitta bellii*  
*Kinosternon sonoriensis*  
*Pseudemys scripta gaigeae*

##### Eastern

*Deirochelys reticularia*  
*Malaclemmys pileata littoralis*  
*Terrapene carolina major*  
*Pseudemys floridana hoyi*  
*Pseudemys floridana mobilensis*

#### DISCUSSION

*Importance of Escarpment Area.* In Table 1 are summarized the effects of the Balcones Escarpment on the distribution of various groups of amphibians and reptiles. The alligator (*Alligator mississippiensis* Daudin) occurs in Texas, but is not included in the table. About 56 per cent of the 216 amphibians and reptiles recorded for Texas reach or cross the escarpment. The Balcones region forms a natural range limit for 77 per cent of the 122 forms involved in the area. The variation among the different groups is from 67 to 100 per cent. Least influenced are turtles, and next in order of increasing influence are snakes, anurans, lizards, and finally salamanders. The variation is correlated with habitat preferences, physiological tolerances, and life history requirements.

It is of interest to note that the percentage of forms reaching the escarpment area varies relatively little (50 to 67) among the five groups studied. In other words (1) these groups are nearly equally well distributed over the central part of the state, and (2) the proportion of the fauna of the state occurring in the escarpment area is sufficiently large to be of significance as a basis for zoogeographic considerations of general import.

Somewhat similarly indicative is the comparison of the combined counts for all groups of (1) eastern plus northern species and (2) western species, which include most (all but 22) forms in the state with the deletion of the endemic forms. A ratio of virtually 1-1 is obtained (76 western, 79 eastern plus northern).

*Limiting Effect of Escarpment.* Of those reaching the escarpment, 77 per cent are limited in range at that point, or, as a corollary, 23 per cent occur on both sides. If this were accepted as a general indication, it may be stated that a 75 per cent degree of differentiation in the herpetofauna at any boundary would be required to match the category into which the escarpment bound-

TABLE 1

## SUMMARY OF EFFECTS OF ESCARPMENT

Group	Anurans	Salamanders	Lizards	Snakes	Turtles	Total
I. Number in state	38	18	44	93	23	216
II. Number not reaching escarpment	19	6	21	40	8	94
III. Number reaching escarpment	19	12	23	53	15	122
IV. Number limited	14	12	21	37	10	94
V. Endemics	2 (10%)	5 (42%)	1 (4%)	2 (4%)	1 (7%)	11 (9%)
VI. Number on both sides	5	0	2	16	5	28
VII. Per cent reaching escarpment	50	67	52	57	65	56
VIII. Per cent limited	74	100	95	70	67	77
IX. Per cent, VI of III	26 (11)	0	5	30	33 (7)	23
X. Ratio, W to E + N in II + IV	0.5 (9-17)	0.1 (1-11)	3.1 (28-9)	1.2 (34-29)	0.3 (4-13)	1.0 (76-79)
XI. Ratio, W to E, in IV	0.7 (5-7)	0.2 (1-5)	1.4 (11-8)	0.8 (14-17)	0.1 (1-8)	0.7 (32-45)
XII. Ratio, W to E + N, in II	0.4 (4-10)	0.0 (0-6)	17.0 (17-1)	1.7 (20-12)	0.6 (3-5)	1.3 (44-34)

ary may be placed. However, a different figure has been arrived at for the birds (56 per cent), a deficiency which may be attributed to the aerial habits of most birds.

Such wide differences in range limitation of the several groups belonging to only two classes of vertebrates at such a conspicuous natural boundary as that formed by the Balcones Escarpment indicate that consideration of all groups plant and animal is extremely important in designating the biotic areas of this region of Texas.

It is not improbable that, when adequately known in detail, some "index" group, as for instance the amphibians and reptiles, could be used at least tentatively to establish boundaries of biotic areas. A fairly precise table of correlation should be worked out in advance between the per cent of limitation (or differentiation) and rank of the biotic area.

The degree of endemism (autochthony is not distinguished) is not high. Not all cases counted are clear-cut, and those that are admitted average only 9 per cent for all groups, with a variation of from 42 per cent in salamanders to 4 per cent in snakes and lizards. Endemism apparently varies with extent of isolation from competition, not with degree of adaptability. Salamanders find conditions conducive to isolation more than any other group, inhabiting there, as they do, numerous well-separated artesian drainage systems. Frogs and turtles with their dependence upon humid conditions, find a lesser degree of opportunity for isolation. Least opportunity for isolation is offered to snakes and lizards, and in these the least endemism is found.

**Differentiation Contours** The percentage of limitation at the escarpment exhibited by reptiles and amphibians is relatively high. It is obvious that the percentage decreases to the north and south, as the sharpness of the boundary lessens. This phenomenon—distinctness of a barrier in one area, and a gradual lessening of importance away from that area suggests the value of plotting degree of differentiation of a given fauna by lines spaced at a given interval as for instance 10 per cent. The lines resulting would resemble physiographic contour lines, with a close approach of the lines in areas of great differentiation, a spreading in areas of little differentiation as in the Great Plains. Such contours, if constructed, would aid greatly in the evaluation of biotic provinces.

Differentiation contours in theory would have the advantage of combining the features of the "biotic province" and the "life zone" concepts, each of which by itself is deficient as a measure of natural areas. And again, differentiation contours lend themselves not only to W-E differentiation but also N-S or any other directional line of differentiation.

The difficulties of constructing such contours on a large scale as for the entire United States are such that the idea may be impractical for a number of years; however, it is at least a possible means of natural mapping that deserves consideration.

*Degree of Tolerance.* Data furnished by the distribution of the herpetofauna in the Balcones Escarpment area offer a means of establishing at least roughly the degree of tolerance by the various groups concerned of the varying conditions occurring in the area (see Table 2). To be considered separately are the average tolerance of the species and the tolerance of the group as a whole. These might well be considered jointly, save for the fact that while numerous species of a given group may overlap a given boundary like the escarpment, the group as a whole is concentrated on one side or the other of the same boundary.

TABLE 2  
SUMMARY OF POSSIBLE INTERPRETATIONS

Criteria	Indications		Group Allocation
	Tolerance	Status of Group	
Unbalanced X-Y ratio	group—low	minor,	
High percentage of boundary overlap	species—high	ascendent	
Balanced X-Y ratio	group—high	dominant,	Serpentes
High percentage of boundary overlap	species—high	ascendent	
Unbalanced X-Y ratio	group—low	minor,	Anura Caudata Lacertilia Chelonia
Low percentage of boundary overlap	species—low	descendent	
Balanced X-Y ratio	group—high	dominant,	
Low percentage of boundary overlap	species—low	descendent	

Degree of tolerance is expressed by two sets of data: by the degree of overlap at the escarpment, and by the ratio of western forms to eastern forms. For these analyses it is preferable to consider only forms reaching the plateau, disregarding the entire state fauna, in order to eliminate a possible source of error introduced by artificial location of political boundaries. The extent of overlap appears to be an index to degree of tolerance of individual species, while the West-East ratio (hereafter called x-y ratio for the sake of brevity and to permit application to situations in which different cardinal directions may be involved from the simple West-East ratio that is important in this particular case) appears to be an index to degree of tolerance in the group as a whole.

Species of salamanders, with the least crossing and with next to the most unbalanced x-y ratio, are obviously the least tolerant of all, individually and as a group, of the varied conditions exhibited in the escarpment area. Species of lizards are next, with very little overlap and a distinctly unbalanced x-y ratio (in favor of the western group); obviously lizards are relatively intolerant of mesic conditions. In anurans a rather high per cent of crossing occurs, and the x-y ratio is less unbalanced; as a group they have been able to adjust more successfully to all the peculiar variations in the area of the escarpment than either salamanders or lizards. The data on crossing are undoubtedly distorted, however, due to the fact that anurans, like turtles, can live under essentially mesic conditions near or in streams that penetrate far into areas of xeric conditions. Although five forms are listed as crossing over, three demand mesic conditions, and therefore the actual per cent of crossing might better be regarded as nearer 10 than **26**. There is still a higher percentage of crossing in snakes, and a more closely balanced x-y ratio; snakes are indicated as the most tolerant of all groups considered. Turtles, with an apparently high degree of crossing, are incorrectly allocated by these data, since all but one of those crossing the area are aquatic and thus are not subject to the same limiting factors that affect the distribution of the other groups. The tolerance of turtles is better indicated by the extremely unbalanced x-y ratio, which would mark them as even less tolerant than salamanders.

*Status of Groups.* Among the five groups comprising the herpetofauna of the escarpment area only two combinations of crossing percentages and x-y ratios are represented: (1) a high percentage of crossing combined with a balanced x-y ratio (Serpentes only), and (2) a low percentage of crossing combined with an unbalanced x-y ratio (Anura, Caudata, Lacertilia, Chelonina). There are, however, theoretically two other combinations possible: (3) a high percentage of crossing with an unbalanced x-y ratio, and (4) a low percentage of crossing with a balanced x-y ratio.

It is not unreasonable to suppose that groups with a low tolerance as a whole (an unbalanced x-y ratio) are *minor* groups, as compared with *dominant* groups which possess a high tolerance as a whole. In like fashion, groups whose species possess high tolerance may be considered as *ascendent*, inasmuch as their species are capable, at the present time, of investigating and inhabiting a wide variety of niches; these are distinct from *descendent* groups whose species have become adapted morphologically for life under certain set conditions and do not enter territories with widely different conditions.

Accordingly, snakes may be regarded as now a dominant and ascendent group, vigorous in an evolutionary sense; all other groups here considered



(anurans, salamanders, lizards, turtles) are minor and decadent groups probably receding in importance on the evolutionary scale.

#### CONCLUSIONS

1. The herpetofauna in the area of the Balcones Escarpment is 77 per cent limited, with a variation from 67 to 100 per cent in the five groups considered.

2. This per cent of limitation is greater than for birds (56 per cent).

3. "Biotic area" boundaries, indicated therefore by such widely varying degrees of limitation, should be considered on the basis of all groups—**plant** and animal—concerned.

4. "Index" groups to biotic areas should, however, have some significance when correlations are established between degree of limitation and rank of area.

5. An autochthony of 9 per cent is present and may be correlated approximately with a limitation of an entire fauna of about 77 per cent.

6. Differentiation (or Limitation) Contours may be more useful than either Life Zones or Biotic Provinces in mapping the biota of large areas.

7A. Degree of tolerance in gross terms can be interpreted for individual species and for groups as a whole, from data on percentage of crossing and on x-y ratio, respectively.

7B. Likewise the evolutionary status (i.e., whether dominant or minor, ascendent or descendent) of groups can be determined also from crossing percentages and from x-y ratio.

a. Much crossing of an unbalanced fauna indicates a high specific and low group tolerance, and occurs in minor ascendent groups.

b. Much crossing of a balanced fauna indicates a high specific and high group tolerance, and occurs in dominant, ascendent groups.

c. Little crossing of an unbalanced fauna indicates a low specific and low group tolerance, and occurs in minor, descendent groups.

d. Little crossing of a balanced fauna indicates a low specific and high group tolerance, and occurs in dominant, descendent groups.

8. Snakes are regarded as a dominant and ascendent group.

9. Anurans, salamanders, lizards, and turtles are all regarded as minor, descendent groups.

## LITERATURE CITED

Buechner, Helmut Karl

- 1944 The range vegetation of Kerr County, Texas, in relation to livestock and white-tailed deer. *Amer. Midi. Nat.*, **vol. 31, no. 3**, p. 697-743.
- 1946 Birds of Kerr County, Texas. *Trans. Kansas Acad. Sci.*, **vol. 49, no. 3**, p. 357-364, fig. 1.

Bishop, S. C.

- 1943 Handbook of salamanders of the United States, Canada, and Lower California. Comstock Publ. Co., Ithaca, N. Y., **xiv + 555 p., 144 fig., 56 maps.**

Pope, C. H.

- 1939 Turtles of the United States and Canada. Alfred A. Knopf, New York, **xviii + 343 p., 99 fig.**

Schmidt, K. P. and Davis, D. Dwight

- 1941 Field book of snakes of the United States and Canada. G. P. Putnam's Sons, New York, **xiii + 365 p., 103 fig., 34 pl.**

Smith, H. M.

- 1946 Handbook of lizards. Comstock Publ. Co., Ithaca, N. Y., **xxi + 557 p., 136 figs., 135 pl., 41 maps.**

Tharp, B. C.

- 1939 The vegetation of Texas. *Texas Acad. Sci. Publ. Nat. Hist., Nontech. Ser.*, **xvi + 74 p., 1 pl., 17 fig.**

Wright, A. H. and Wright, A. A.

- 1938 Amphibians of Texas. *Trans. Texas Acad. Sci.*, **vol. 21, p. 1-38, pl. 1-3, text fig. 1-6.**
- 1942 Handbook of frogs and toads. Comstock Publ. Co., Ithaca, N. Y., **xi + 286 p., 88 pl.**